

Notes for Miscellaneous Lectures

Leonid A. Levin*
Boston University†

Abstract

Here I share a few notes I used in various course lectures, talks, etc. Some may be just calculations that in the textbooks are more complicated, scattered, or less specific; others may be simple observations I found useful or curious.

1 Nemirovski Estimate of Mean of Arbitrary Distributions with Bounded Variance

The popular Chernoff bounds assume severe restrictions on distribution: it must be cut-off, or vanish exponentially, etc. In [Nemirovsky Yudin]¹ an equally simple bound uses no conditions at all beyond independence and known bound on variance. It is not widely used because it is not explained anywhere with an explicit tight computation. I offer this version:

Assume independent variables $X_i(\omega)$ with the same unknown mean m and known lower bounds B_i^2 on inverses $1/v_i$ of their variance. We estimate m as $M(\omega)$ with probability $P(\pm(M-m) \geq \varepsilon) \stackrel{\text{def}}{=} p^\pm < 2^{-k}$ for k close to $\sum (B_i \varepsilon)^2 / 12$. First, we normalize X_i to set $\varepsilon = 1$, spread them into n groups, and take in each group j its B_i^2 -weighted mean $x_j(\omega)$.

The inverse variance bounds b_j^2 for x_j are additive; we grow groups to assure $b_j > 2$ and to increase the sum k of *heights* $h_j \stackrel{\text{def}}{=} \log_2 \frac{b_j + b_j^{-1}}{2}$. (The best $h/b^2 > 1/12$ comes with $b^2 \approx 6$.)²

For $s \subset [1, n]$, let $b_s \stackrel{\text{def}}{=} \prod_{j \in s} b_j$. Let $L \stackrel{\text{def}}{=} \cup_t L_t$ consist of *light* s , whose largest superset s' with $b_{s'}^2 < b_{[1, n]}$ has $\|s\| + t$ elements. As $s \in L_t$ do not include each other, $\|L_t\| \leq \binom{n}{\lceil n/2 \rceil} < 2^n \sqrt{2/(\pi n)}$, by Sperner's theorem, and since $n! = (n/e)^n \sqrt{2\pi n + \theta_n}$, $\pi/3 < \theta_n \leq e^2 - 2\pi$.

Our M is the $(\log b_j)$ -weighted median of x_j . Let $S^\pm(\omega) \stackrel{\text{def}}{=} \{j : \pm(x_j - m) < 1\}$. Then $\pm(M(\omega) - m) \geq 1$ means $S^\pm(\omega) \in L$. By Chebyshev's inequality, $p_j^\pm \stackrel{\text{def}}{=} P(j \notin S^\pm) \leq 1/(b_j^2 + 1)$. We assume $p_j^\pm = 1/(b_j^2 + 1)$: the general case follows by so modifying the distribution without changing m , b_j , or decreasing p^+ (respectively p^-). If $s \in L_t$, $S^\pm(\omega) = s$ has probability

$$p_s^\pm = b_s^2 / \prod_{j \leq n} (b_j^2 + 1) < 4^{-t} \prod_{j \leq n} b_j / (b_j^2 + 1) = 4^{-t} 2^{-(k+n)}. \text{ So,}$$

$$p^+ + p^- \leq \sum_{t \geq 0} \sum_{s \in L_t} (p_s^+ + p_s^-) \leq 2 \left(\sum_{t \geq 0} 4^{-t} \right) 2^{-(k+n)} 2^n \sqrt{2/(\pi n)} < 2^{-k} \sqrt{5/n}. \blacksquare$$

*Supported by NSF grant 0311411.

†Computer Science department, 111 Cummington St., Boston, MA 02215, USA

¹A.S.Nemirovsky, D.B.Yudin. *Problem Complexity and Method Efficiency in Optimization*. Wiley, 1983.

²Giving up tightness, the rest may be simplified: assure $b_j \geq b^{\text{def}} \sqrt{2} + 1$ and replace b_j, h_j with $b, 1/2$.

2 Leftover Hash Lemma

The following Lemma is often useful to convert a stream of symbols with absolutely unknown (except for a lower bound on its entropy) distribution into a source of perfectly uniform random bits $b \in Z_2 = \{0, 1\}$.

The version I give is close to that in [HILL]³, though some aspects are closer to that from [GL]⁴. Unlike [GL], I do not restrict hash functions to be linear and do not guarantee polynomial reductions, i.e. I forfeit the case when the unpredictability of the source has computational, rather than truly random, nature. However, like [GL], I restrict hash functions only in probability of collisions, not requiring pairwise uniform distribution.

Let G be a probability distribution on Z_2^n with Renyi entropy $-\log \sum_x G^2(x) \geq m$. Let $f_h(x) \in Z_2^k$, $h \in Z_2^t$, $x \in Z_2^n$ be a hash function family in the sense that for each $x, y \neq x$ the fraction of h with $f_h(x) = f_h(y)$ is $\leq 2^{-k} + 2^{-m}$. Let U^t be the uniform probability distribution on Z_2^t and $s = m - k - 1$. Consider a distribution $P(h, a) = 2^{-t} G(f_h^{-1}(a))$ generated by identity and f from $U^t \otimes G$. Let $\mathbf{L}_1(P, Q) = \sum_z |P(z) - Q(z)|$ be the $\mathbf{L}_1$ distance between distributions P and $Q = U^i$, $i = t + k$. It never exceeds their $\mathbf{L}_2$ distance

$$\mathbf{L}_2(P, Q) = \sqrt{2^i \sum_z (P(z) - Q(z))^2}.$$

Lemma 1 (Leftover Hash Lemma)

$$\mathbf{L}_1(P, U^i) \leq \mathbf{L}_2(P, U^i) < 2^{-s/2}.$$

Note that h must be uniformly distributed but can be reused for many different x . These x need to be independent only of h , not of each other as long as they have $\geq m$ entropy in the distribution conditional on all their predecessors.

Proof.

$$\begin{aligned} (\mathbf{L}_2(P, U))^2 &= 2^i \sum_{h,a} P(h, a)^2 + 2^i \sum_z (2^{-2i} - 2P(z)2^{-i}) = 2^i \sum_{h,a} P(h, a)^2 - 1 \\ &= -1 + 2^i \sum_{x,y} G(x)G(y)2^{-2t} \sum_a \|\{h: f_h(x) = f_h(y) = a\}\| \\ &= -1 + 2^{k-t} \sum_{x,y} G(x)G(y) \|\{h: f_h(x) = f_h(y)\}\| \\ &= -1 + 2^{k-t} \left(\sum_x G(x)^2 2^t + \sum_{x,y \neq x} G(x)G(y) \|\{h: f_h(x) = f_h(y)\}\| \right) \\ &\leq -1 + 2^k 2^{-m} + 2^{k-t} (1 - 2^{-m}) 2^t (2^{-k} + 2^{-m}) < 2^{-s}. \end{aligned}$$

■

³Johan Hastad, Russell Impagliazzo, Leonid A. Levin, Michael Luby.

A Pseudorandom Generator from any One-way Function. Section 4.5. *SICOMP* **28**(4):1364-1396, 1999.

⁴Oded Goldreich, Leonid A. Levin. A Hard-core Predicate for any One-way Function. Sec.5. *STOC* 1989.

3 Disputed Ballots and Poll Instabilities

Here is another curious example of advantages of quadratic norms.

The ever-vigilant struggle of major parties for the heart of the median voter makes many elections quite tight. Add the Electoral College system of the US Presidential elections and the history may hang on a small number of ballots in one state. The problem is not in the randomness of the outcome. In fact, chance brings a sort of fair power sharing unplagued with indecision: either party wins sometimes, but the country always has only one leader. If a close race must be settled by dice, so be it. But the dice must be trusty and immune to manipulation!

Alas, this is not what our systems assure. Of course, old democratic traditions help avoiding outrages endangering younger democracies, such as Ukraine. Yet, we do not want parties to compete on tricks that may decide the elections: appointing partisan election officials or judges, easing voter access in sympathetic districts, etc. Better to make the randomness of the outcome explicit, giving each candidate a chance depending on his/her share of the vote. It is easy to implement the lottery in an infallible way, the issue is how its chance should depend on the share of votes.

In contrast to the present one, the system should avoid any big jump from a small change in the number of votes. Yet, chance should not be proportional to the share of votes. Otherwise each voter may vote for himself, rendering election of a random person. The present system encourages voters to consolidate around candidates acceptable to many others. The ‘jumpless’ system should preserve this feature. This can be done by using a non-linear function: say the chance in the post-poll lottery be proportional to the *squared* number of votes. In other words, a voter has one vote per each person he agrees with.⁵ Consider for instance an 8-way race where the percents of votes are 60, 25, 10, 1, 1, 1, 1, 1. The leader’s chance will be $5/6$, his main rival’s $1/7$, the third party candidate’s $1/43$ and the combined chance of the five ‘protest’ runners $1/866$.

This system would force major parties to determine the most popular candidate via some sort of primaries, and will almost exclude marginal runners. However it would have no discontinuity rendering any small change in the vote distribution irrelevant. The system would preserve an element of chance, but would be resistant to manipulation.

⁵The dependence of lottery odds on the share of votes may be sharper. Yet, it must be smooth to minimize the effects of manipulation. Even (trusty) noise alone, e.g., discarding a randomly chosen half of the votes, can “smooth” the system a little.

4 Proofs in Three Envelopes

Below is a slightly simplified account of Zero-knowledge proofs that were developed in [Goldwasser Micali Rackoff,⁶ Goldreich Micali Wigderson,⁷ Shamir⁸]. I wrote this account under the influence of Manuel Blum's constructions⁹ during several conversations with him when I visited him in 1986.

Consider an undirected graph: $g \in v^2$, $(a, b) \in g$ iff $(b, a) \in g$. Its coloring is a mapping $C : v \rightarrow \{1, 2, 3\}$, s.t. each edge $(a, b) \in g$ has distinct colors: $C(a) \neq C(b)$. Since 3-colorability is NP-complete, any mathematical statement can be reduced in polynomial time to a statement of graph colorability, so that any proof of either statement can be transformed in polynomial time into the proof of the other. We consider only graphs composed of 3 isomorphic connected components. Any coloring of such graph can be made *balanced*, i.e. such that the nodes of each degree are equally spread between the 3 colors. We consider only such *balanced* colorings.

The Prover (P) uses a random string ω to generate random enumerations $p : v \rightarrow v$ of nodes and $q : g \rightarrow g$ of edges. Then P makes three envelopes: $E_1(g, \omega)$, $E_2(g, \omega)$, $E_3(g, \omega, C)$. E_1 contains p and the mapping of reciprocal edges: $q(a, b) \rightarrow q(b, a)$. E_2 contains the mapping of edges to their source nodes: $q(a, b) \rightarrow p(a)$ and E_3 contains their coloring $q(a, b) \rightarrow C(a)$. The verifier (V) then chooses any two of the envelopes and checks their consistency.

If the envelopes do not represent a correct coloring then some two of them are obviously inconsistent with the graph or with each other. It is also easy to see that the joint probability distribution of any two envelopes does not depend on (balanced) coloring and can be trivially generated from the graph alone. E_1, E_2 do not mention C at all. E_1, E_3 contain just p and an unrelated to it balanced mapping of permuted edges to their colors. E_2, E_3 contain the permutation and colors of nodes and also maps to them their permuted outgoing edges with unspecified destination.

So the Prover gives away no information besides the validity of his proof, while the verifier has a 1/3 chance to catch him if the proof is incorrect. Repeating the game k times with independent ω decreases the chance of fake proofs to remain un-exposed to $(2/3)^k$. Of course, for implementing such game one needs something like cryptography to commit the Prover to the content of the envelopes, without revealing it before he learns the verifier's choice.

4.1 Graph non-isomorphism

A simple protocol of [Goldreich Micali Wigderson] shows in zero knowledge that an *isomorphism* of two graphs g_1, g_2 is known. P first sends V a random permutation h of g_1 . Then V chooses at random $i \in \{1, 2\}$ and P sends V the isomorphism of h to g_i . Non-isomorphism [Goldreich Micali Wigderson] has almost as simple protocol. Let g be the graph whose connected components are g_1 and g_2 . V sends P a random permutation h of g **and** proves in zero-knowledge that (s)he knows an isomorphism of h to g . Then P tells V if the permutation maps the two components of g onto themselves or onto each other.

⁶Shafi Goldwasser, Silvio Micali, Charles Rackoff. The Knowledge Complexity of Interactive Proofs. *SICOMP* **18**:186-208, 1989. Earlier version in *STOC*-1985.

⁷Oded Goldreich, Silvio Micali, Avi Wigderson. Proofs that Yield Nothing but their Validity or All Languages in NP have Zero-Knowledge Proof Systems. *JACM*, **38**(1):691-729, 1991. Earlier in *FOCS*-1986.

⁸Adi Shamir. Zero-Knowledge Proof for Knapsacks. Cited in: Joe Kilian, Silvio Micali, Rafail Ostrovsky. Minimum Resource Zero-Knowledge Proof. [FOCS] 1989, pp. 474-479.

⁹Manuel Blum. How to prove a theorem so no one else can claim it. *Proc. 1986 International Congress of Math.* Also, Personal Communication. Berkeley, 1986.